

(1995 年)

- 1) Kitakaze M, Hori M, Morioka T, Minamino T, Takashima S, Okazaki Y, Node K, Komamura K, Iwakura K, Itoh T, et al.

Alpha 1-adrenoceptor activation increases ecto-5'-nucleotidase activity and adenosine release in rat cardiomyocytes by activating protein kinase C.

Circulation. 1995 Apr 15;91(8):2226-34. PMID: 7697853.

- 2) Mangrulkar R. S, Ono M, Ishikawa M, Takashima S, Klagsbrun M, Nowak R. A.

Isolation and Characterization of Heparin-Binding Growth- Factors in Human Leiomyomas and Normal Myometrium.

Biol Reprod. 1995 Sep;53(3):636-46. PMID: 7578688.

- 3) Morioka T, Kitakaze M, Minamino T, Takashima S, Node K, Okazaki Y, Sato H, Shinozaki Y, Chujo M, Mori H, et al.

Downward shift of coronary pressure-flow relationship following a brief period of ischemia in dogs.

Am J Physiol. 1995 Oct;269(4 Pt 2):H1237-45. PMID: 7485554.

(1994 年)

- 1) Kitakaze M, Hori M, Minamino T, Takashima S, Komamura K, Node K, Kurihara T, Morioka T, Sato H, Inoue M, et al.

Evidence for deactivation of both ectosolic and cytosolic 5'-nucleotidase by adenosine A1 receptor activation in the rat cardiomyocytes.

J Clin Invest. 1994 Dec;94(6):2451-6. PMID: 7989602; PMCID: PMC330077.

- 2) Kitakaze M, Hori M, Morioka T, Minamino T, Takashima S, Sato H, Shinozaki Y, Chujo M, Mori H, Inoue M,

et al.

Alpha 1-adrenoceptor activation mediates the infarct size-limiting effect of ischemic preconditioning through augmentation of 5'- nucleotidase activity.

J Clin Invest. 1994 May;93(5):2197-205. PMID:8182151; PMCID: PMC294363.

- 3) Kitakaze M, Hori M, Morioka T, Minamino T, Takashima S, Sato H, Shinozaki Y, Chujo M, Mori H, Inoue M,

et al.

Infarct size-limiting effect of ischemic preconditioning is blunted by inhibition of 5'-nucleotidase activity and attenuation of adenosine release.

Circulation. 1994 Mar;89(3):1237-46. PMID: 8124812.

- 4) Hori M, Kitakaze M, Takashima S, Morioka T, Sato H, Minamino T, Node K, Komamura K, Inoue M, Kamada T.

AICA riboside improves myocardial ischemia in coronary microembolization in dogs.

Am J Physiol. 1994 Oct;267(4 Pt 2):H1483-95. PMID: 7943395.

(1993 年)

- 1) Kitakaze M, Hori M, Takashima S, Sato H, Inoue M, Kamada T.
Ischemic preconditioning increases adenosine release and 5'-nucleotidase activity during myocardial ischemia and reperfusion in dogs. Implications for myocardial salvage.
Circulation. 1993 Jan;87(1):208-15. Erratum in: *Circulation* 1993 May;87(5):1775. *Circulation* 1993 Jun;87(6):2070. PMID: 8419009.
- 2) Sato H, Hori M, Kitakaze M, Iwai K, Takashima S, Kurihara H, Inoue M, Kamada T.
Reperfusion after brief ischemia disrupts the microtubule network in canine hearts.
Circ Res. 1993 Feb;72(2):361-75. PMID: 8418989.
- 3) Kitakaze M, Hori M, Takashima S, Iwai K, Sato H, Inoue M, Kitabatake A, Kamada T.
Superoxide dismutase enhances ischemia-induced reactive hyperemic flow and adenosine release in dogs. A role of 5'-nucleotidase activity.
Circulation. 1993 Mar;87(3):982-95. PMID: 8443917.
- 4) Kitakaze M, Hori M, Morioka T, Takashima S, Minamino T, Sato H, Inoue M, Kamada T.
Attenuation of ecto-5'-nucleotidase activity and adenosine release in activated human polymorphonuclear leukocytes.
Circ Res. 1993 Sep;73(3):524-33. PMID: 8348695.
- 5) Hori M, Kitakaze M, Morioka T, Minamino T, Takashima S, Sato H, Node K, Shinozaki Y, Chujo M, Mori H.
An increase in adenosine release contributes to the infarct size limiting effect of ischemic preconditioning.
Jpn Circ J. 1993;57 Suppl 4:1330-2. Japanese. PMID: 7966979.
- 6) Kitakaze M, Hori M, Takashima S, Morioka T, Minamino T, Sato H, Okazaki Y, Inoue M, Kamada T.
Superoxide dismutase enhances both adenosine release and coronary hyperemic flow through protection of 5'-nucleotidase against its degradation during reperfusion following ischemia in dogs.
Biorheology. 1993 Sep-Dec;30(5-6):359-70. PMID: 8186402.

(1992 年)

- 1) Sato H, Hori M, Kitakaze M, Takashima S, Inoue M, Kitabatake A, Kamada T.
Endogenous adenosine blunts beta-adrenoceptor-mediated inotropic response in hypoperfused canine myocardium.
Circulation. 1992 Apr;85(4):1594-603. PMID: 1313344.

(1991 年)

- 1) Hori M, Kitakaze M, Sato H, Takashima S, Iwakura K, Inoue M, Kitabatake A, Kamada T.
Staged reperfusion attenuates myocardial stunning in dogs. Role of transient acidosis during early reperfusion.
Circulation. 1991 Nov;84(5):2135-45. PMID: 1657451.
- 2) Kitakaze M, Hori M, Sato H, Takashima S, Inoue M, Kitabatake A, Kamada T.
Endogenous adenosine inhibits platelet aggregation during myocardial ischemia in dogs.
Circ Res. 1991 Nov;69(5):1402-8. PMID: 1657446.