

(2002 年)

- 1) Asakura M, Kitakaze M, Takashima S, Liao Y, Ishikura F, Yoshinaka T, Ohmoto H, Node K, Yoshino K, Ishiguro H, Asanuma H, Sanada S, Matsumura Y, Takeda H, Beppu S, Tada M, Hori M, Higashiyama S. Cardiac hypertrophy is inhibited by antagonism of ADAM12 processing of HB-EGF: metalloproteinase inhibitors as a new therapy.
Nat Med. 2002 Jan;8(1):35-40. PMID: 11786904.
- 2) Takashima S, Kitakaze M, Asakura M, Asanuma H, Sanada S, Tashiro F, Niwa H, Miyazaki Ji J, Hirota S, Kitamura Y, Kitsukawa T, Fujisawa H, Klagsbrun M, Hori M. Targeting of both mouse neuropilin-1 and neuropilin-2 genes severely impairs developmental yolk sac and embryonic angiogenesis.
Proc Natl Acad Sci U S A. 2002 Mar 19;99(6):3657-62. Epub 2002 Mar 12. PMID: 11891274; PMCID: PMC122579.
- 3) Soker S, Miao H. Q, Nomi M, Takashima S, Klagsbrun M. VEGF(165) mediates formation of complexes containing VEGFR-2 and neuropilin-1 that enhance VEGF(165)-receptor binding.
J Cell Biochem. 2002;85(2):357-68. PMID: 11948691.
- 4) Liao Y, Ishikura F, Beppu S, Asakura M, Takashima S, Asanuma H, Sanada S, Kim J, Ogita H, Kuzuya T, Node K, Kitakaze M, Hori M. Echocardiographic assessment of LV hypertrophy and function in aortic-banded mice: necropsy validation.
Am J Physiol Heart Circ Physiol. 2002 May;282(5):H1703-8. PMID: 11959634.
- 5) Kitakaze M, Asanuma H, Funaya H, Node K, Takashima S, Sanada S, Asakura M, Ogita H, Kim J, Hori M. Angiotensin-converting enzyme inhibitors and angiotensin II receptor blockers synergistically increase coronary blood flow in canine ischemic myocardium: role of bradykinin.
J Am Coll Cardiol. 2002 Jul 3;40(1):162-6. PMID: 12103271.
- 6) Sanada S, Node K, Asanuma H, Ogita H, Takashima S, Minamino T, Asakura M, Liao Y, Ogai A, Kim J, Hori M, Kitakaze M. Opening of the adenosine triphosphate-sensitive potassium channel attenuates cardiac remodeling induced by long-term inhibition of nitric oxide synthesis: role of 70-kDa S6 kinase and extracellular signal-regulated kinase.
J Am Coll Cardiol. 2002 Sep 4;40(5):991-7. PMID: 12225728.
- 7) Asakura M, Kitakaze M, Sakata Y, Asanuma H, Sanada S, Kim J, Ogida H, Liao Y, Node K, Takashima S, Tada M, Hori M. Adenosine-induced cardiac gene expression of ischemic murine hearts revealed by cDNA array hybridization.
Circ J. 2002 Jan;66(1):93-6. PMID: 11999673.
- 8) Ogita H, Node K, Asanuma H, Sanada S, Takashima S, Asakura M, Kitakaze M, Hori M. Amelioration of ischemia- and reperfusion-induced myocardial injury by the selective estrogen receptor modulator, raloxifene, in the canine heart.
J Cardiol. 2002 Jan;39(1):55-6. Japanese. PMID: 11828800.

(2001 年)

- 1) Sanada S, Kitakaze M, Papst PJ, Asanuma H, Node K, Takashima S, Asakura M, Ogita H, Liao Y, Sakata Y, Ogai A, Fukushima T, Yamada J, Shinozaki Y, Kuzuya T, Mori H, Terada N, Hori M. Cardioprotective effect afforded by transient exposure to phosphodiesterase III inhibitors: the role of protein kinase A and p38 mitogen-activated protein kinase. *Circulation*. 2001 Aug 7;104(6):705-10. PMID: 11489779.
- 2) Kitakaze M, Node K, Takashima S, Asanuma H, Asakura M, Sanada S, Shinozaki Y, Mori H, Sato H, Kuzuya T, Hori M. Role of cellular acidosis in production of nitric oxide in canine ischemic myocardium. *J Mol Cell Cardiol*. 2001 Sep;33(9):1727-37. PMID: 11549351.
- 3) Sanada S, Kitakaze M, Papst PJ, Hatanaka K, Asanuma H, Aki T, Shinozaki Y, Ogita H, Node K, Takashima S, Asakura M, Yamada J, Fukushima T, Ogai A, Kuzuya T, Mori H, Terada N, Yoshida K, Hori M. Role of phasic dynamism of p38 mitogen-activated protein kinase activation in ischemic preconditioning of the canine heart. *Circ Res*. 2001 Feb 2;88(2):175-80. PMID: 11157669.
- 4) Sanada S, Kitakaze M, Node K, Takashima S, Ogai A, Asanuma H, Sakata Y, Asakura M, Ogita H, Liao Y, Fukushima T, Yamada J, Minamino T, Kuzuya T, Hori M. Differential subcellular actions of ACE inhibitors and AT(1) receptor antagonists on cardiac remodeling induced by chronic inhibition of NO synthesis in rats. *Hypertension*. 2001 Sep;38(3):404-11. PMID: 11566913.
- 5) Asanuma H, Kitakaze M, Node K, Takashima S, Sakata Y, Asakura M, Sanada S, Shinozaki Y, Mori H, Tada M, Kuzuya T, Hori M. Benidipine, a long-acting Ca channel blocker, limits infarct size via bradykinin- and NO-dependent mechanisms in canine hearts. *Cardiovasc Drugs Ther*. 2001;15(3):225-31. PMID: 11713890.
- 6) Asanuma H, Kitakaze M, Funaya H, Takashima S, Minamino T, Node K, Sakata Y, Asakura M, Sanada S, Shinozaki Y, Mori H, Kuzuya T, Tada M, Hori M. Nifedipine limits infarct size via NO-dependent mechanisms in dogs. *Basic Res Cardiol*. 2001 Sep;96(5):497-505. PMID: 11605997.
- 7) Fukushima N, Ohtake S, Sawa Y, Nishimura M, Matsumiya G, Takashima S, Hori M, Shirakura R, Matsuda H. Follow-up on heart transplant after left ventricular assist system bridge. *Asian Cardiovascular and Thoracic Annals*. 9:312-314.
- 8) Sanada S, Kitakaze M, Asanuma H, Harada K, Ogita H, Node K, Takashima S, Sakata Y, Asakura M, Shinozaki Y, Mori H, Kuzuya T, Hori M. Role of mitochondrial and sarcolemmal K(ATP) channels in ischemic preconditioning of the canine heart. *Am J Physiol Heart Circ Physiol*. 2001 Jan;280(1):H256-63. PMID: 11123240.
- 9) Kitakaze M, Asakura M, Sakata Y, Asanuma H, Sanada S, Kuzuya T, Miyazaki J, Takashima S, Hori M. cDNA array hybridization reveals cardiac gene expression in acute ischemic murine hearts. *Cardiovasc Drugs Ther*. 2001 Mar;15(2):125-30. PMID: 11669405.