

(2004 年)

- 1) Asano Y, Takashima S, Asakura M, Shintani Y, Liao Y, Minamino T, Asanuma H, Sanada S, Kim J, Ogai A, Fukushima T, Oikawa Y, Okazaki Y, Kaneda Y, Sato M, Miyazaki J, Kitamura S, Tomoike H, Kitakaze M, Hori M.  
Lamr1 functional retroposon causes right ventricular dysplasia in mice.  
*Nat Genet.* 2004 Feb;36(2):123-30. Epub 2004 Jan 18. PMID: 14730304.
- 2) Liao Y, Asakura M, Takashima S, Ogai A, Asano Y, Shintani Y, Minamino T, Asanuma H, Sanada S, Kim J, Kitamura S, Tomoike H, Hori M, Kitakaze M.  
Celiprolol, a vasodilatory beta-blocker, inhibits pressure overload-induced cardiac hypertrophy and prevents the transition to heart failure via nitric oxide-dependent mechanisms in mice.  
*Circulation.* 2004 Aug 10;110(6):692-9. Epub 2004 Jul 19. PMID: 15262839.
- 3) Sanada S, Asanuma H, Minamino T, Node K, Takashima S, Okuda H, Shinozaki Y, Ogai A, Fujita M, Hirata A, Kim J, Asano Y, Mori H, Tomoike H, Kitamura S, Hori M, Kitakaze M.  
Optimal windows of statin use for immediate infarct limitation: 5'-nucleotidase as another downstream molecule of phosphatidylinositol 3-kinase.  
*Circulation.* 2004 Oct 12;110(15):2143-9. Epub 2004 Sep 27. PMID: 15451788.
- 4) Shintani Y, Node K, Asanuma H, Sanada S, Takashima S, Asano Y, Liao Y, Fujita M, Hirata A, Shinozaki Y, Fukushima T, Nagamachi Y, Okuda H, Kim J, Tomoike H, Hori M, Kitakaze M.  
Opening of Ca<sup>2+</sup>-activated K<sup>+</sup> channels is involved in ischemic preconditioning in canine hearts.  
*J Mol Cell Cardiol.* 2004 Dec;37(6):1213-8. PMID: 15572051.
- 5) Asanuma H, Sanada S, Ogai A, Minamino T, Takashima S, Asakura M, Ogita H, Shinozaki Y, Mori H, Node K, Tomoike H, Hori M, Kitakaze M.  
Methotrexate and MX-68, a new derivative of methotrexate, limit infarct size via adenosine-dependent mechanisms in canine hearts.  
*J Cardiovasc Pharmacol.* 2004 Apr;43(4):574-9. PMID: 15085069.
- 6) Sanada S, Asanuma H, Tsukamoto O, Minamino T, Node K, Takashima S, Fukushima T, Ogai A, Shinozaki Y, Fujita M, Hirata A, Okuda H, Shimokawa H, Tomoike H, Hori M, Kitakaze M.  
Protein kinase A as another mediator of ischemic preconditioning independent of protein kinase C.  
*Circulation.* 2004 Jul 6;110(1):51-7. Epub 2004 Jun 21. PMID: 15210595.
- 7) Okada K, Minamino T, Tsukamoto Y, Liao Y, Tsukamoto O, Takashima S, Hirata A, Fujita M, Nagamachi Y, Nakatani T, Yutani C, Ozawa K, Ogawa S, Tomoike H, Hori M, Kitakaze M.  
Prolonged endoplasmic reticulum stress in hypertrophic and failing heart after aortic constriction: possible contribution of endoplasmic reticulum stress to cardiac myocyte apoptosis.  
*Circulation.* 2004 Aug 10;110(6):705-12. Epub 2004 Aug 2. PMID: 15289376.
- 8) Ogita H, Node K, Asanuma H, Sanada S, Kim J, Takashima S, Minamino T, Hori M, Kitakaze M.  
Raloxifene improves coronary perfusion, cardiac contractility, and myocardial metabolism in the ischemic heart: role of phosphatidylinositol 3-kinase/Akt pathway.  
*J Cardiovasc Pharmacol.* 2004 Jun;43(6):821-9. PMID: 15167276.
- 9) Ogita H, Node K, Liao Y, Ishikura F, Beppu S, Asanuma H, Sanada S, Takashima S, Minamino T, Hori M, Kitakaze M.  
Raloxifene prevents cardiac hypertrophy and dysfunction in pressure-overloaded mice.  
*Hypertension.* 2004 Feb;43(2):237-42. Epub 2003 Dec 15. PMID: 14676219.

10) Asanuma H, Minamino T, Sanada S, Takashima S, Ogita H, Ogai A, Asakura M, Liao Y, Asano Y, Shintani Y, Kim J, Shinozaki Y, Mori H, Node K, Kitamura S, Tomoike H, Hori M, Kitakaze M. Beta-adrenoceptor blocker carvedilol provides cardioprotection via an adenosine-dependent mechanism in ischemic canine hearts.

*Circulation*. 2004 Jun 8;109(22):2773-9. Epub 2004 May 17. PMID: 15148268.

(2003 年)

1) Takashima S, Iwamoto R, Yamazaki S, Asakura M, Hasuwa H, Miyado K, Adachi S, Kitakaze M, Hashimoto K, Raab G, Nanba D, Higashiyama S, Hori M, Klagsbrun M, Mekada E.

Heparin-binding EGF-like growth factor and ErbB signaling is essential for heart function.

*Proc Natl Acad Sci U S A*. 2003 Mar 18;100(6):3221-6. Epub 2003 Mar 5. PMID: 12621152; PMCID: PMC152273.

2) Yamazaki S, Iwamoto R, Saeki K, Asakura M, Takashima S, Yamazaki A, Kimura R, Mizushima H, Moribe H, Higashiyama S, Endoh M, Kaneda Y, Takagi S, Itami S, Takeda N, Yamada G, Mekada E.

Mice with defects in HB-EGF ectodomain shedding show severe developmental abnormalities.

*J Cell Biol*. 2003 Nov 10;163(3):469-75. Epub 2003 Nov 3. PMID: 14597776; PMCID: PMC2173637.

3) Liao Y, Takashima S, Asano Y, Asakura M, Ogai A, Shintani Y, Minamino T, Asanuma H, Sanada S, Kim J, Ogita H, Tomoike H, Hori M, Kitakaze M.

Activation of adenosine A1 receptor attenuates cardiac hypertrophy and prevents heart failure in murine left ventricular pressure-overload model.

*Circ Res*. 2003 Oct 17;93(8):759-66. Epub 2003 Sep 11. PMID: 12970111.

4) Asakura M, Takashima S, Asano Y, Honma T, Asanuma H, Sanada S, Shintani Y, Liao Y, Kim J, Ogita H, Node K, Minamino T, Yorikane R, Agai A, Kitamura S, Tomoike H, Hori M, Kitakaze M.

Canine DNA array as a potential tool for combining physiology and molecular biology.

*Circ J*. 2003 Sep;67(9):788-92. PMID: 12939556.

5) Asanuma H, Node K, Minamino T, Sanada S, Takashima S, Ueda Y, Sakata Y, Asakura M, Kim J, Ogita H, Tada M, Hori M, Kitakaze M.

Celiprolol increases coronary blood flow and reduces severity of myocardial ischemia via nitric oxide release.

*J Cardiovasc Pharmacol*. 2003 Apr;41(4):499-505. PMID: 12658050.

6) Ogita H, Node K, Asanuma H, Sanada S, Takashima S, Minamino T, Soma M, Kim J, Hori M, Kitakaze M. Eicosapentaenoic acid reduces myocardial injury induced by ischemia and reperfusion in rabbit hearts.

*J Cardiovasc Pharmacol*. 2003 Jun;41(6):964-9. PMID: 12775977.

7) Sanada S, Node K, Minamino T, Takashima S, Ogai A, Asanuma H, Ogita H, Liao Y, Asakura M, Kim J, Hori M, Kitakaze M.

Long-acting Ca<sup>2+</sup> blockers prevent myocardial remodeling induced by chronic NO inhibition in rats.

*Hypertension*. 2003 Apr;41(4):963-7. Epub 2003 Mar 10. PMID: 12629037.

8) Asano Y, Kim J, Ogai A, Takashima S, Shintani Y, Minamino T, Kitamura S, Tomoike H, Hori M, Kitakaze, M.

A calcium channel blocker activates both ecto-5'(-)-nucleotidase and NO synthase in HUVEC.

*Biochem Biophys Res Commun*. 2003 Nov 21; 311(3):625-8. PMID: 14623316.